

Lunar-driven control of climate and Barents Sea eco-systems¹

Harald Yndestad²

Introduction

Herring periods and cod periods along the Norwegian coast have been known for more than 1000 years. Periods of growth in the fisheries, have formed the basis for settlement, industrialization, economic growth, and wealth. Periods, when the fish disappeared, led to emigration, hunger, and poverty. Over the years, one has questioned whether good years, or bad years, were accidental, or ruled by higher powers. The fish stock grew during the 1940s. After 1945, a new fishing fleet was built, which had good years in the 1950s and 1960s. In the 1970s, the herring stock and the cod stock disappeared. Questions were then asked about possible causes. Was there a lack of scientific management, overfishing, or was there something unknown phenomenon in nature, which led to the fish stocks disappearing. This presentation is a summary of a doctoral dissertation in which this topic was studied.

The 18.6 year cycle

Gravity between the earth, the sun, and the moon, cause the earth's axis to change direction, in a period of 18.6 years. The earth's axis nutation period introduces a standing lunar nodal tidal wave, between the pole and the equator, with a period of 18.6 years. The tidal wave has a maximum amplitude at the poles and the equator and a node at about 35 degrees in the orbital plane.





Figure 1. The Earth axis and the lunar nodal tide wave.

¹ The talk can be seen here: <https://www.youtube.com/watch?v=EN1WwV9TZ7Y> (Recorded by Yngvar Engebretsen).

² Submitted Dec. 20, 2021. Accepted Dec. 30, 2021. DOI: <https://doi.org/10.53234/scc202112//222>

The tidal wave of 18.6 years leads to a vertical mixing of between cold and warm temperature layers. The result of vertical mixing is that the surface water has a periodic change of 18.6 years, which follows the ocean currents northwards, and affects the climate. Over time, a set of temperature periods of 18.6 years is produced, which gives periodic spectrum of $[1, 2, 3, 4 \dots] \cdot 18.6$ years, which gives climate periods up to 75 and 223 years. A wavelet spectrum analysis of data series has identified periods of $[1/2, 1, 4] \cdot 18.6$ years in the inflow of North Atlantic water to the Norwegian Sea, Barents Sea, the NAO index, land surface temperature and rainfall along the Norwegian coastline. Arctic ice extent from 1570, has periodic variations in periods of $[1, 4, 12] \cdot 18.6$ years.



(Yndestad et al. 2008)

Figure 2. The North Atlantic water temperature cycle.


The effect of the cycle on the fisheries

A study of Barents Sea ecosystems shows that green algae, plankton, herring, capelin, and cod follow a periodic variation of $[1/2, 1, 3, 4] \cdot 18.6$ years. The capelin stock recruits in periods of $(18.6/2)/3$ years, which optimizes the capelin stock over a period of $18.6/2$ years. The cod stock recruits in periods of $18.6/3$ years, which optimizes the stock's growth over 18.6 years. The cod stock period of 18.6 years, at the same time follows the temperature variations of $3 \cdot 18.6$ years and $4 \cdot 18.6$ years, which has an average period of $(3 + 4)18.6/2 = 65$ years. Herring periods follow cod stock periods. The explanation for good fishery periods and bad fishery periods, is recruitment, growth, and mortality, adapted to sea temperature variations of $[1/2, 1, 2, 3, 4, \dots] \cdot 18.6$ years. The explanation for the biomass collapse, is allocation of catch quotas, which does not follow the sea temperature variations of $[1/2, 1, 2, 3, 4, \dots] \cdot 18.6$ years.

See also www.climateclock.no.

Keywords: Lunar nodal tide; Lunar forced climate variability; Marine ecosystem variability; Eco system resonance; Biomass collapse


Phase-locked to $18.6/2=9.3$ yr NAW temp period



(Yndestad and Stene, 2001)

Figure 3. . The 9.3 years cycle and its influence on the capelin.

Period- and phase-locked to lunar-driven NAW temperature



(Yndestad, 1999, 2009)

Figure 4. The 18.6 years cycle and its influence on the cod.

References

Yndestad, H 1996, **System Dynamics of North Arctic Cod**, *The 84'th international ICES Annual Science Conference, Hydrography Committee, Iceland, October 1996.*

- Yndestad, H 1997. **Systems Dynamics in the Fisheries of Northeast Arctic Cod.** *15th International System Dynamics Conference (ISDC '97), August 1997, Istanbul.*
- Yndestad, H 2000. **The predestined fate. The Earth nutation as a forced oscillator on management of Northeast Arctic cod.** *The 18th International Conference of The System Dynamics Society. Aug. 6-10, 2000, Bergen, Norway.*
- Yndestad, H 2001a. **Earth nutation influence on Northeast Arctic cod management.** ICES Journal of Marine Science, 58, 799-805.
- Yndestad H and Stene A 2002. **Systems Dynamics of Barents Sea Capelin.** ICES Journal of Marine Science, 59, 1155-1166.
- Yndestad H 2002. **The Code of Norwegian spring spawning herring Long-term cycles.** *ICES Annual Science Conference, Oct. 2002, Copenhagen.*
- Yndestad H 2003a. **A Lunar nodal spectrum in Arctic time series.** *ICES Annual Science Conference, Sept. 2003. Tallinn. ICES CM 2003/T.*
- Yndestad H, Turrell W R and Ozhigin V 2004. **Temporal linkages between the Faro-Shetland time series and the Kola section time series.** *ICES Annual Science Conference, Vigo. Sept. 2004. Theme Session M. Regime Shifts in the North Atlantic Ocean: Coherent or Chaotic?*
- Yndestad H. 2004. **The cause of Barents Sea biomass dynamics.** Journal of Marine Systems.
- Yndestad, H. Dr.philos. Thesis. 2004:132. *The Lunar nodal cycle influence on the Barents Sea.* Norges teknisk-naturvitenskapelige universitet, NTNU. Trondheim.
- Yndestad, H. 2006. **The Arctic Ocean as a coupled oscillating system to the forced 18.6 yr lunar nodal cycle.** *20 Years of Nonlinear Dynamics in Geosciences. American Meteorological Society & European Geosciences Union, June 11-16, 2006, Rhodes, Greece.*
- Yndestad, H. 2006. **The influence of the lunar nodal cycle on Arctic climate.** ICES Journal of Marine Science, 63:3, 401–420. <http://doi.org/10.1016/j.icesjms.2005.07.015>
- Yndestad, H 2006. **Possible Lunar tide effects on climate and the ecosystem variability in the Nordic Seas and the Barents Sea.** *ICES annual conference. Session ICES CM 2006/C: Climatic variability in the ICES area 2000-2005 in relation to previous decades: physical and biological consequences 19-23 sept. 2006, Maastricht, Netherland.*
- Yndestad, H 2007. **Long tides influence on the climate dynamics and the ecosystem dynamics in the Barents Sea.** *Symposium on Ecosystem Dynamics in the Norwegian Sea and the Barents Sea. Theme session: Climatic effects on food webs. Tromsø. Norway 12-15 March 2007.*
- Yndestad, H 2008. **The Barents Sea ecosystem dynamics as a coupled oscillator to long tides.** *Annual Science conference 22-26 Sept. 2008, Theme Session Coupled physical and biological models: parameterization, validation and application. ICES CM 2008/L:01*
- Yndestad, H, Turrell W R and Ozhigin, V 2008. **Lunar nodal tide effects on variability of sea level, temperature, and salinity in the Faroe-Shetland Channel and the Barents Sea.** Deep-Sea Research I: Oceanographic Research Papers, 55:10, 1201–1217. <http://doi.org/10.1016/j.dsr.2008.06.003>
- Yndestad H 2009. **The influence of long tides on ecosystem dynamics in the Barents Sea.** Deep-Sea Research II. (2009) 2108-2116. Deep-Sea Research II, 56, 2108-2116. <http://doi.org/10.1016/j.dsr2.2008.11.022>
- Yndestad, H 2021. **Barents Sea Ice Edge Position Variability 1579-2020**, Report no TN210214, NTNU-Ålesund. May 2021. <http://doi.org/10.13140/RG.2.2.16122.41928>